

**Instructions for the Extemporaneous Compounding of:
Pyrimethamine Suspension 2mg/ml**

Batch Number: _____

Formulation:

Pyrimethamine (Daraprim®) 25 mg	2 tab (50mg)	4 tabs (100mg)
Keltrol 0.4%w/v <u>IQ</u>	25mL	50mL

Formula	Quantity	Batch No.	Expiry Date	Manufacturer	Prepared By	Checked By
Pyrimethamine 25mg tablet (Daraprim®)				GlaxoSmithKline		
Keltrol 0.4%w/v <u>IQ</u>				Victoria Pharmaceuticals		

Instructions for Preparation:

1. Print the labels and leave to one side.
2. Using a sterile alcohol wipe, clean the mortar and pestle. Allow to dry.
3. Count the required number of tablets. **CHECK.**
4. Grind the tablets to a fine powder in the mortar for 10mins.
5. Use a sufficient portion of Keltrol 0.4%w/v to make a smooth paste.
6. Slowly add the Keltrol 0.4%w/v in increments. Mix well after each addition until the pyrimethamine is fully suspended.
7. Transfer to a clean, dry graduated cylinder.
8. Make up to the final volume with Keltrol 0.4%w/v. **CHECK.**
9. Transfer to suitable sterile bottle. Cap and label appropriately.
10. Leave to one side for checking. **FINAL CHECK**

EXPIRY: 7 days when stored at room temperature.

Manufactured date: _____

Expiry date: _____

Prepared By: _____

Checked By: _____

Additional Labels Required:

DISPENSED LABEL:

Disclaimer: CAUTION: The information on The OLCHC Pharmacy Department Web Site has been developed specifically for OLCHC. It is provided for informational purposes for qualified health care professionals. This material is not intended as a substitute for consulting qualified health care professionals. Patient circumstances will vary and some information may have become outdated as a result of more recent medical developments. The Pharmacy Department at Our Lady's Children's Hospital, Crumlin makes no representation, express or implied, with regard to the accuracy of the information contained in this publication and cannot accept any legal responsibility for any errors or omissions that may be made.